AD-A119 655

TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY

MACROCYCLES CONTAINING TIN. TWO SYNTHESES OF 1.1.6.6.11.11.16.1--ETC(U)

SEP 82 M NEWCOMB, Y AZUMA, A R COURTNEY

N00014-79-C-0584

NL

END

SEND

HO 392

Onc.

(1.2)

Unclassified

SECURITY CLASSIF'CATION OF THIS PAGE (When Lete Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
Technical Report 3 $ADA1196$	3. RECIPIENT'S CATALOG NUMBER
Macrocycles Containing Tin. Two Syntheses of 1,1,6,6,11,11,16,16-Octaphenyl-1,6,11,16-tetrastannacycloeicosane and a Synthesis of 1,1,6,6-Tetraphenyl-1,6-distannacyclodecane	5. TYPE OF REPORT & PERIOD COVERSON Technical Report 6. PERFORMING ORG. REPORT NUMBER
Martin Newcomb, Yutaka Azuma, Arleen R. Courtney	N00014-79-C-0584
PERFORMING ORGANIZATION NAME AND ADDRESS Texas A&M University Department of Chemistry College Station, TX 77843	NR 053-714
Office of Naval Research Department of the Navy Arlington, Virginia 22217 14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office)	12. REPORT DATE Sept. 22, 1982 13. NUMBER OF PAGES 10 15. SECURITY CLASS. (of this report)
16. DISTRIBUTION STATEMENT (of this Report)	Unclassified 15a. DECLASSIFICATION/DOWNGRACING SCHEDULE

Approved for public release; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)



18. SUPPLEMENTARY NOTES

H

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Polystanna macrocycle Synthesis Purification

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

Synthetic techniques and methods for analysis and purification of macrocycles containing tin atoms are exemplified by the title syntheses. Highlights of the syntheses include a highly selective conversion of 1,4-bis-(triphenyl-stannyl) butane to 1,4-bis-(bromodiphenylstannyl) butane and a two component macrocyclization which proceeds in 43% yield to give the 20-membered ring product. Analytical HPLC and preparative chromatography methods for characterization and purification of the macrocycles and their synthetic intermediates

DD 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE 5/N 0102-014-6601 | Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

LLUMITY CLASSIFICATION OF THIS PAGE(When Data Entered)

and analogs are described. Other syntheses which show the generality of the methods are discussed.

Accession For

NTIS GRAAI
DTIC TAB
Unamounced
Justification

By
Distribution/
Availability Codes
Avail and/or
Dist Special

Unclassified

The Property of the Control of the C

SECURITY CLASSIFICATION OF THIS PAGE(When Date Entered)

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0584

Task No. NR 053-714
TECHNICAL REPORT NO. 3

Macrocycles Containing Tin.

Two Syntheses of

1,1,6,6,11,11,16,16-Octaphenyl-1,6,11,16-tetrastannacycloeicosane

and a Synthesis of

1,1,6,6-Tetrapheny1-1,6-distannacyclodecane

by

Martin Newcomb, Yutaka Azuma, and Arleen R. Courtney

Texas A&M University Department of Chemistry College Station, TX 77843

Sept. 22, 1982

Reproduction in whole or in part is permitted for any purpose of the United States Government

- * This document has been approved for public release and sale; its distribution is unlimited
- * This statement should also appear in Item 10 of Document Control Data DD Form 1473. Copies of form available from cognizant contract administrator.

Macrocycles Containing Tin. Two Syntheses of 1,1,6,6,11,11,16,16-Octaphenyl-1,6,11,16-tetrastannacycloeicosane and a Synthesis of 1,1,6,6-Tetraphenyl-1,6-distannacyclodecane

Martin Newcomb*¹, Yutaka Azuma, and Arleen R. Courtney

Department of Chemistry, Texas A&M University, College Station, Texas 77843

Summary: The title syntheses are described. The synthetic and purification and analytical methods employed are of general utility for the preparation and functionalization of members of this class of compounds.

Macrocyclic, polydentate cation-complexing ligands (crown ethers, cryptands, etc.) have enjoyed a broad and useful chemistry during the past decade and a half, but their counterparts, anion-complexing macrocycles have received relatively little attention. Recent advances in anion complexation, including structural selectivity, by macrocyclic polyammonium ligands suggest that this will be a fertile area of stidy. Reasoning that polystanna macrocycles may be appropriately substituted to give Lewis acid complexing ligands which are direct analogs of crown ethers or cryptands, we have developed synthetic procedures for this class of compounds represented by the title compound (1). Two approaches (Scheme) have yielded the target 20-membered ring compound. The synthetic methods and purification techniques described herein are generally applicable for this class of compounds and also furnish tin function lized macrocycles. We are unaware of other neutral macrocycles with the potential for anion complexation.

All reactions were run in argon or nitrogen atmospheres. Treatment of the diGrignard reagent from 1,4-dibromobutane with triphenylstanryl chloride in tetrahydrofuran (THF) gave 1,4-bis-(triphenylstannyl)butane 3a,b (2, mp 148.5-149 °C) in 77% isolated yield after recrystallization from hexane--dichloromothane (2:1, v:v). Unacceptably low selectivity was observed in several attempted conversions of 2 to 1,4-bis-(bromodiphenylstannyl)butane (3) with various reagents and conditions; a second phenyl group was readily replaced giving the dibromophenylstannyl moiety. However, treatment of 2 with 2.1 molar equivalents of hydrogen bromide in dry dichloromethane at -78 °C followed by slow warming to room temperature gave, after recrystallization (dry ether), 3^{3a} in 75% yield (mp 88-90 °C). The dibromide 3 was treated with excess 4-chlorobuty1magnesium bromide in THF at -10 $^{
m O}{
m C}$ for 1.5 h followed by warming to room temperature (7 h) to give 1,14-dichloro-5,5,10,10-tetraphenyl-5,10-distannatetradecane $(1)^{3a}$ which was purified by reverse phase chromatography (C-18, methanol elution) in 62% yield (oil).

The two component macrocyclization was accomplished at high diltation. The dibromide 3 was added to excess lithium metal in THF to give the dilithium reagent 5 (total base = 63% of theory). A THF solution of 5 (diluted to 0.037 M) was added slowly (2 h) to a THF solution of 4 (1.0 molar equivalent, 0.02 M) at 0 $^{\circ}$ C followed by warming to room temperature (12 h). After a conventional work-up, the crude product was purified by reverse phase chromatography (C-18) with THF--acetonitrile elution (1:3, v:v) to give the desired macrocycle 1^3 in 43% isolated yield (mp 107.5-108 $^{\circ}$ C from hexane--ether, 3:1).

Alternatively, macrocycle 2 was also obtained in lower yield from a high dilution, four component macrocyclization reaction. Thus, the diGrignard reagent from 1,4-dibromobutane in THF (0.07 M) was added over 2 h to one molar equivalent of dibromide 3 in THF (0.034 M) at 0 °C. After 12 h at room temperature, the reaction was quenched and worked up. The crude products were purified by preparative reverse phase chromatography as above to give, after recrystallization, the desired macrocycle 1 in 163 yield and 1,1,6,6-tetraphenyl-1,6-distannacyclodecane 3a,5 (6, mp 114-114.5 °C, lit. 4 mp 96-98 °C) in 10% yield. Compound 6 has been prepared by a different route. 4

The macrocycles 1 and 6 may be analyzed readily by analytical HPLC (reverse phase, C-18) with methanol elution; the 10-membered ring compound 6 elutes before 1. However, for preparative chromatography, the solubilities of 1 and 6 in methanol are inconveniently low. Thus, chromatography with the mixed solvent system THF+-acetonitrile was developed. We found that analytical HPLC on 10 μ (pherisorb ODS columns correlated well with preparative chromatography on 40 μ ODS supplied by J. T. Baker Co.; the analytical phase retained material about 1.5 times as long as the preparative phase in terms of column volumes.

The procedures described above are generally useful for the preparation and functionalization of other tin containing macrocycles. For example, the reactions of the 6-, 8-, and 10-carbon analogs of 3 with the corresponding chain length a,w-diGrignard reagents gave the 14-, 18-, and 22-membered ring analogs of the distanna compound 6, respectively, as well as low yields of the 28-, 36-, and 44-membered ring analogs of tetrastanna compound 1, respectively. All separations were accomplished by reverse phase chromatography. Further, when the selective bromination procedure was applied to macrocycle 6, we obtained 1,6-dibromo-1,6-diphenyl-1,6-distannacyclodecane; 3a this reaction exemplifies a critically important functionalization of the macrocycles. Finally, preparative reverse phase chromatography of the intermediate tin bromides is also possible on a C-18 column with THF--acetonitrile elution if dry solvents are used.

As in any macrocyclization reaction, the high dilution methods we used required careful technique. However, in our minds, the key steps to obtaining macrocycle 1 were the selective bromination of 2 and our development of a preparative chromatography method. The two component macrocyclization route and reverse phase preparative chromatography permit the synthesis of 1 in gram batches. With the methods at hand we plan to prepare and functionalize several members of this class of compounds and explore their application in anion coordination chemistry.

Acknowledgement: This work was supported by the Office of Naval Research.

References and Notes

- (1) Camille and Henry Dreyfus Teacher-Scholar, 1980-1985.
- (2) Dietrich, B.; Hosseini, M. W.; Lehn, J. M.; Sessions, R. B.
 <u>J. Am. Chem. Soc.</u> 1981, 103, 1282; Hosseini, M. W.; Lehn, J. M.
 <u>J. Am. Chem. Soc.</u> 1982, 104, 3525; and references in each.
- (3) (a) The compound was characterized by ¹H NMR spectroscopy; (b) the compound was characterized by ¹H-decoupled ¹³C NMR spectroscopy and by osmometric molecular weight determination; (c) a satisfactory elemental analysis (±0.4% for C, H) vas obtained.
- (4) Davies, A. G.; Tse, M. -W.; Kennedy, J. D.; McFarlane, W.; Pyne,
 G. S.; Ladd, M. F. C.; Povey, D. C. J. Chem. Soc., Chem. Commun.
 1978, 791.

Scheme

Supplementary Information for Review Purposes

Osmometric Molecular Weight Determinations.

Compound	MW Calcd	MW Found
Ę	755	734
1	1315	1341
6	657	654

Elemental Analysis of Compound χ .

 $\mathrm{C}_{64}\mathrm{H}_{72}\mathrm{Sn}_4\colon\,\text{\%C}$ calcd 58.38, found 58.06; $\,\text{\%H}$ calcd 5.52, found 5.43.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No.		No.
	Copies		Copies
Office of Naval Research		Naval Ocean Systems Center	
Attn: Code 413		Attn: Mr. Joe McCartney	
800 North Quincy Street		San Diego, California 92152	1
Arlington, Virginia 22217	2	•	
-		Naval Weapons Center	
ONR Pasadena Detachment		Attn: Dr. A. B. Amster,	
Attn: Dr. R. J. Marcus		Chemistry Division	
1030 East Green Street		China Lake, California 93555	1
Pasadena, California 91106	1		
		Naval Civil Engineering Laboratory	
Commander, Vaval Air Systems Command]	Attn: Dr. R. W. Drisko	
Attn: Code 310C (H. Rosenwasser)		Port Hueneme, California 93401	1
Department of the Navy			
Washington, D.C. 20360	1	Dean William Tolles	
		Naval Postgraduate School	
Defense Technical Information Center	-	Monterey, California 93940	1
Building 5, Cameron Station			
Alexandria, Virginia 22314	12	Scientific Advisor	
		Commandant of the Marine Corps	
Dr. Fred Saalfeld		(Code RD-1)	
Chemistry Division, Code 6100		Washington, D.C. 20380	1
Naval Research Laboratory			
Washington, D.C. 20375	1	Naval Ship Research and Development Center	
U.S. Army Research Office		Attn: Dr. G. Bosmajian, Applied	
Attn: CRD-AA-IP		Chemistry Division	
P. O. Box 12211		Annapolis, Maryland 21401	1
Research Triangle Park, N.C. 27709	1		
•		Mr. John Boyle	
Mr. Vincent Schaper		Materials Branch	
DTNSRDC Code 2803		Naval Ship Engineering Center	
Annapolis, Maryland 21402	1	Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center		Mr. A. M. Anzalone	
Attn: Dr. S. Yamamoro		Administrative Librarian	
Marine Sciences Division		PLASTEC/ARRADCOM	
San Diego, Californi: 91232	1	Bldg 3401	
		Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	No.		No.
	Copies		Copies
Dr. M. F. Hawthorne		Dr. T. Marks	
Department of Chemistry		Department of Chemistry	
University of California		Northwestern University	
Los Angeles, California 90024	1	Evanston, Illinois 60201	1
Dr. D. B. Brown		Dr. J. Zuckerman	
Department of Chemistry		Department of Chemistry	
University of Vermont		University of Oklahoma	
Burlington, Vermont 05401	1	Norman, Oklahoma 73019	1
Dr. D. Venezky		Professor O. T. Beachley	
Chemistry Division		Department of Chemistry	
Naval Research Laboratory		State University of New York	
Code 6130		Buffalo, New York 14214	1
Washington, D.C. 20375	1		
5 ,		Professor K. M. Nicholas	
Dr. John E. Jensen		Department of Chemistry	
Hughes Research Laboratory		Boston College	
3011 Malibu Canyon Road		Chestnut Hill, Massachusetts 02167	1
Malibu, California 90265			
•		Professor R. Neilson	
Dr. A. Cowley		Department of Chemistry	
Department of Chemis:ry		Texas Christian University	
University of Texas		Fort Worth, Texas 76129	1
Austin, Texas 78712	1	. •	
		Professor M. Newcomb	
Dr. W. Hatfield		Texas A&M University	
Department of Chemistry		Legartment of Chemistry	
University of North Carolina		College Station, Texas 77843	1
Chapel Hill, North Carolina 27514	1		
		Professor Richard Eisenberg	
Dr. M. H. Chislolm		Department of Chemistry	
Department of Chemis:ry		University of Rochester	
Indiana University		Rochester, New York 14627	1
Bloomington, Indiana 47401	1		
		Professor R. D. Archer	
		University of Massachusetts	
		Chemistry Department	
		Amherst, Massachusetts 01003	

DATE